

ARO 19618.1-C

ig(ig) Stereochemistry of Reactions Involving

Cyclic Peroxides

Final Repart / // /

Med A. Porter s

Professor of Chemistry

14 | 25 September 1931

U.S. Army Research Office

Proposal #15610-C

Duke University

Duke University

(13)11:

15 Non 116 1

D

DIST

Approved for public releases Distribution Unlimited

REPORT LOCUMENTATION	PAGE	READ INSTRUCTIONS SEFORE COMPLETING FORM
REPORT NUMBER	2. GOVT ACCESSION NO. 2. PEG	
	AD-4105999	
GTCE (m. ! Skude)		POLECTIONE SPERIOD COVERE
Stereochemistry of Reactions Inv Cyclic Peroxides	olving Find	al Report by 1978-30 April 81
7 Found		TRACT OR GRANT NUMBER(A)
Ned A. Porter	NSF	CHE 77-26088 Propusal 15618-C
PERFORMING ONE AND A COUNTRAKE ALL MODRESS	10. Ph	GGRAN SEEMENT PROJECT, YAS
Duke University Durham NC		EAR VORK UNIT HUMBERS
CONTROLLING OFFICE NAME AND AUDRESS	12. RE	PORT DATE
U. S. Army Research Office		5 September 81
Post Office Box 12211 Research Tricarle Fack, NC 277		HUER OF PAGES
PONITORING AGENCY HAME & ADDRESSAL allient		CURLLY CLASS. (of this report)
	Ţ,	clasaifică
V	15r. 0	ECLASSIFICATION DOVIGENOUS
	5	CHEDULE
Approved for public release; di		Chenge E
Approved for public release; di	stribution unlimited.	
Approved for public release; di	stribution unlimited.	
Approved for public release; di	In Eleck 20, II different from Report ngs contained in this trued as an official	report are those of t
Approved for public release; di DISTRIBUTION STATEMENT (of the abstract enforced NA B. SUPPLEMENTARY NOTES The view, opinions, and/or find author(s) and should not be cons	ngs contained in this trued as an official inless so designated b	report are those of t
Approved for public release; di DISTRIBUTION STATEMENT (of the abeliant unforce) NA Supplementary Notes The view, opinions, and/or find author(s) and should not be consposition, policy, or decision, to	in Flock 20, If different from Report In Struck 20, If different from Report Ings contained in this trued as an official Inless so designated b	report are those of t Department of the Army y other documentation.
Approved for public release; di DISTRIBUTION STATEMENT (of the abstract entered NA Supplementary Notes The view, opinions, and/or find author(s) and should not be consposition, policy, or decision, to key words (Continue on reverse alde If necessary & Autoxidation, cyclic peroxides,	ngs contained in this trued as an official inless so designated by polymer degradation,	report are those of t Department of the Army y other documentation.
Approved for public release; di DISTRIBUTION STATEMENT (of the abstract entered MA Supplementary Notes The view, opinions, and/or find author(s) and should not be consposition, policy, or decision, to key words (Continue on reverse alde If necessary and Autoxidation, cyclic peroxides,	ngs contained in this trued as an official inless so designated by polymer degradation,	report are those of t Department of the Army y other documentation.
Approved for public release; di DISTRIBUTION STATEMENT (of the abstract entered NA Supplementary Notes The view, opinions, and/or find author(s) and should not be consposition, policy, or decision, to key words (Continue on reverse alde If necessary & Autoxidation, cyclic peroxides,	ngs contained in this trued as an official inless so designated by polymer degradation,	report are those of t Department of the Army y other documentation.
Approved for public release; di DISTRIBUTION STATEMENT (of the abstract entered MA B. SUPPLEMENTARY NOTES The view, opinions, and/or find author(s) and should not be consposition, policy, or decision, to be position, policy, or decision, to be authority with the constant of the con	ngs contained in this trued as an official inless so designated by polymer degradation,	report are those of t Department of the Army y other documentation.

20. Abstract

Several aspects of chemistry related to autoxidation, peroxy free radicals, and peroxides have been explored. Methods for preparation of complex unsaturated hydroperoxides have been developed and peroxy radicals derived from these hydroperoxies were studied as models for polyunsaturated fatty acid and polyolefin (i.e., polybutadiene and polyisoprene) autoxidation. We have established that peroxy radical cyclization and bicyclization occurs in models of polyunsaturated fatty acid oxidation and a new type of cyclization, serial cyclization, has been observed in polybutadiene oxidation models. Kinetic and stereochemical aspects of these cyclization reactions have been explored. The free radical oxidation of polyunsaturated fatty acids has been studied mechanistically. In particular, we believe that the H atom donating ability of the medium in which oxidation occurs is an important factor in determining the partitioning of peroxy free radicals between competitive pathways (i.e., ß scission and cyclization). The results of these studies provide a firm basis for understanding the degradation of materials, foodstuffs, etc.



Acce	ssion For	<u>-</u>
DTIC Unani	GRARI TAB Counced	
By	ribution/	
4	lability Codes	
Dist	Avail and/or Special	


a.) Statement of Problem Studied

The destruction of organic substrates by molecular oxygen is perhaps one of the most important organic reactions. Natural and synthetic polymers, biological materials such as lipids, and for that matter, any organic molecule exposed to the atmosphere may react with ground state molecular oxygen. It is no surprise then, that extensive investigations have focused on the kinetics of organic substrate oxidation by 0_2 . The "hydroperoxide mechanism" for autoxidation which is now generally accepted is outlined below.

1. R-H + 1·
$$\longrightarrow$$
 R· + H-1 Initiation
2. R· + C₂ \longrightarrow RO·O·
3. RO-O· + R-H \longrightarrow ROOH + R·
4. Termination

Despite the numerous studies on autoxidation and inhibition, surprisingly little research has been directed towards product study in autoxidation. It is true that for simple molecules like isopropylbenzene the primary autoxidation product is the tertiary hydroperoxide. However, if the organic substrate has several centers of unsaturation, the simple hydroperoxide product may not be found and in its place, cyclic peroxides are formed. For example, squalene absorbs two moles of oxygen upon autoxidation and one of these oxygen molecules is incorporated as a peroxide, rather than a hydroperoxide group. Although no rigorous structure proof was offered for the squalene oxidation product, it seems likely on the basis of subsequent work that the principal product of squalene autoxidation is one of the cyclic peroxides (1 or 2) formed by the mechanism described below:

For some six years we have been involved in studies related to the mechanism of autoxidation and the synthesis of biologicaly important peroxides. These studies have resuled in the first syntheses of the important biological compound PGH_2 and PGG_2 and have provided mechanistic information about autoxidation of fatty acids, phospholipids, and polymer models. In this final report, two aspects of these studies are discussed, namely, (1) a unified mechanism for lipid autoxidation, and (2) a description of the stereochemical requirements of the S_H^2 reaction.



b.) Summary of the Most Important Results

1) Reversibility of Oxygen Entrapment in Autoxidation:

In the autoxidation of linoleic acid, both trans, cis and trans, trans hydroperoxides are formed. Thus, linoleic acid, 1, leads to four different hydroperoxides, 2a-5a. Two of these, 2a and 4a have the trans, cis geometry while 3a and 5a are trans, trans diene isomers. We have recently published a study of factors controlling the product distribution of 2-5. We find, in short, that the ratio of trans, cis/trans, trans products (2+4/3+5) formed during the initial stages (<2%) of oxidation is dependent on temperature and the concentration of linoleic acid. This ratio (2+4/3+5) varies from 4.2 (neat linoleic acid oxidations at 10° C) to 0.23 (0.24 linoleic acid in benzene at 50° C). The trans, cis/trans, trans ratio is presented in Figure 2 as a function of [L-H] (the concentration of linoleic acid in the oxidation) and temperature. Appropriate controls have been run to insure that the product ratios observed are independent of extent of oxidation (within the 2% oxidation range) and 0_2 pressure.

The essential competition that accounts for the observed $\frac{\text{trans,cis/trans,trans}}{\text{trans,trans}}$ ratio is β scission at peroxy radical conformer 7 (leading ultimately to a trans, trans product) vs. H atom abstraction by radical conformer pair 6/7 to give the trans, cis product.

A kinetic derivation describing this system gives:

$$\frac{\text{trans,cis}}{\text{trans,trans}} = \frac{k_p[L-H]}{k_B(1-\alpha)} + \frac{\alpha}{(1-\alpha)}$$
 Eq. 1

That is, the product ratio depends on [L-H] as is observed in Figure 2 and the valuable kinetic parameters kg and α can be extracted from the data. (α is the partition factor of radical 8 between the trans, cis and trans, trans manifold.) Further, estimates of ΔH_{B}^{\sharp} , the enthalpy of the β scission process, can be obtained from temperature dependence data and ΔH_{B}^{\sharp} is calculated to be in the 16-18 kcal/mole range. We should note that Benson had proposed in a classic paper that peroxy radical β scission is an important reaction in autoxidation. We have simply shown that this β scission process has a profound influence on product stereochemistry. We should also note that linoleic acid is the only good H atom dnor available in this autoxidation system but in reality $k_{\rm P}[{\rm L-H}]$ in Eq. 1 should be replaced by a summation term that includes all H atom donors,

$$\sum_{n=1}^{i} k_{p}^{n} [R_{n}-H] = KP.$$

We have chosen to define this important parameter as KP and it represents the total H atom donating ability of the medium of autoxidation.

2) Stereochemistry of the $S_{\rm H}{\rm i}$ Reaction on Peroxide


Little is known about the stereochemical requirements of the $S_{H}2$ reaction. An $S_{N}2$ type transitin state has been proposed and strong evidence has been presented that suggests that halogen atom attack on cyclopropane carbon occurs with stereochemical inversion. The $S_{H}2$ reaction is apparently extremely sensitive to steric hindrance, and attack on carbon is limited to the strained cyclopropane series. Radical attack on heteroatoms such as 0, N, or S is a much more common event, possibly because the steric hindrance of attack is much less for these dia and tri-valent atoms than it is for substitution at tetravalent carbon. Skell

suggests that "perhaps this type of displacement" (inversion) "will be found to be important for an unhindered substrate".


The S_H i reaction reported here (Scheme 1) offers the possibility of studying the geometric requirements of S_H2 attack on the peroxide linkage since the attacking radical center and the peroxide bond are geometrically fixed by the cyclic peroxide molecular frame. We suggest that the critical geometric parameter for the S_H i reaction is the dihedral angle, ζ , about the 0-C bond between the attacking radical and the leaving oxygen. Further, we propose that for maximum reactivity, this dihedral angle must be 180° . Our evidence in support of this hypothesis has been published in two recent papers, the results of which are summarized here.

A series of β -bromo peroxides were reacted with Bu3Sn-H affording the corresponding alkyl radicals. As an example, the bromide $\underline{9}$ was shown to react as follows:

By carrying out reactions with different concentrations of Bu₃Sn-H, the rate constant for $S_{H}i$ can be determined. These rate constants, $k_{S_{H}}i$, for a series of radicals are shown below.



Rate constants for k_{S_Hi} range generally from $10^4\text{--}10^6$ sec $^{-1}$ for the series of radic radicals $\underline{10\text{--}15}$ with two exceptions, radicals $\underline{13}$ and $\underline{15}$. In both of those species, the attacking radical cannot approach the peroxide bond with $\phi \approx 180^\circ$ because of geometric constraints. In $\underline{13}$, for example, the angle ϕ is 100° at a maximum while in radical $\underline{15}$, the radical center is fixed with $\phi \approx 60^\circ$. Note that the epimeric radical to $\underline{15}$, $\underline{14}$ (which has equitorial substitution and $\phi \approx 180$), has a "normal" k_{S_Hi} .



We conclude from these studies that radical substitution on the peroxide bond does, indeed, follow a colinear approach pathway. The $S_{\rm H}2$ and $S_{\rm N}2$ pathways for reaction on first-row elements are thus apparently similar, with back-side displacement being the rule.

c.) List of Publications

- "The Chemistry of Cyclic Peroxides: The Formation and Decomposition of Prostaglandin Endoperoxide Analogs", Ned A. Porter, John R. Nixon, and Dennis W. Gilmore, Organic Free Radicals, ed., William A. Pryor, American Chemical Society, Washington, D. C., pp. 89-101, 1978.
- "Prostaglandin H₂ Methyl Ester", N. A. Porter, J. D. Byers, R. C. Mebane, D. W. Gilmore, and J. R. Nixon, J. Org. Chem., 43, 2088(1978).
- "Cyclic Peroxides by Intramolecular Peroxymercuration of Unsaturated Hydroperoxides", J. R. Nixon, M. A. Cudd and N. A. Porter, J. Org. Chem., 43, 4048(1978).
- "Separation and Purification of Lecithins by High Pressure Liquid Chromatography, N. A. Porter, R. A. Wolf, and J. R. Nixon, Lipids, 14, 20(1979).
- "Prostaglanding Endoperoxide", N. A. Porter, Free Radicals in Biology, Vol. IV, ed., William A. Pryor, Academic Press, Inc., New York, N.Y., pp. 261-291, 1980.
- "Synthesis of Prostaglandin H ", Ned A. Porter, Jim D. Byers, Katherine M. Holden, and Daniel B. Menzel, J. Amer. Chem. Soc., 101, 4319(1979).
- "The Preparation and Purification of Arachidonic Acid Hydroperoxides of Biological Importance". Ned A. Porter, James Logan, Voula Kontoyiannidou, J. Org. Chem., 44, 3177(1979).
- "The Autoxidation of Arachidnic Acid: Formation of the Proposed SRS-A Intermediate", Ned A. Porter, Richard A. Wolf, Elizabeth M. Yarbro, and Hugo Weenen, Biochem. and Biohys. Res. Comm., 89, 1058(1979).
- "A Fixed Geometry Study of the S 2 Reaction on the Peroxide Bond", Ned A. Porter, M. Amelia Cudd, Richard W. Miller and Andrew T. McPhail, J. Amer. Chem. Soc., 102, 414(1980).
- "Prostaglandin G₂", Ned A. Porter, Jim D. Byers, A. E. Ali, and Thomas A. Eling, J. Amer. Chem. Soc., 102, 1181(1980).
- "The Free Radical Oxidation of Polyunsaturated Lecithins", Ned A. Porter, Richard A. Wolf, and Hugo Weenen, Lipids, 15, 163(1980).
- "A New Route to Lipid Hydroperoxides: Orbital Symmetry Controlled Ring Opening of Vinylcyclopropyl Bromides", Ned A. Porter, David H. Roberts, and Carl B. Ziegler, Jr., J. Amer. Chem. Soc., 102, 5912(1980).
- "The Autoxidation of Poly-Unsaturated Lipids: Factors Controlling the Stereochemistry of Product Hydroperoxides", Ned A. Porter, Bruce A. Weber, Hugo Weenen and Jamil A. Khan, J. Amer. Chem. Soc., 102, 5597(1980).
- "Serial Cyclization of Peroxy Free Radicals: Models for Polyolefin Oxidation", Ned A. Porter, A. Nicholas Rue, and Andrew T. McPhail, J. Amer. Chem. Soc., 102, 5912(1980).

-9-

d.) List of Participating Personnel

a.) List of Participat	tild i el somici	Ph.D. earned
M. Amelia Cudd A. Nicholas Roe James Logan Carl B. Ziegler, Jr. David Roberts	graduate student graduate student graduate student graduate student graduate student	1980 1981